



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.


Larva.—Head rounded, scarcely bilobed, flattened before, erect, free; orange brown, a large, diffuse, black patch on the face of each lobe at the height of the apex of the clypeus, a patch over ocelli and last antennal joint black. Setæ rather coarse, black; width 3.5 mm. Body moderate, cylindrical, uniform, not tapering and not humped. Ground color velvety black; a broad yellowish-white band in the segmental incisures reaching to the line of the feet, joined below the line of the spiracles to a second short band which begins on the middle of the side and extends downward to a little below the line of the feet. A small, anteriorly situated dot dorsally on joints 3, 4, 9–13, sometimes present also on 8 or 7 and 8, but always smaller on these segments. The band in the incisure 2–3 is somewhat broken and the one in 12–13 is narrowed. There is a small dorsal bar representing a band between 13–14, just in front of the anal plate. On joints 5, 6, and 12 the pale bands run a little further ventrally and have a supplementary spot. All feet orange brown, thoracic ones with tips black, the abdominal claspers and tubercle vii black. Cervical shield undifferentiated, black, the anterior edge of joint 2 brown. Anal plate orange brown with a black posterior rim and tubercle each side middle line. Tubercles small, obscure; iv a little below the middle of the spiracle. Setæ rather distinct, black.

DESCRIPTION OF A NEW SESIA.

BY WILLIAM BEUTENMÜLLER.

Sesia palmii, sp. nov.

Head and palpi dark brown; collar mixed with a little white. Antennæ brown black, with a bluish lustre. Thorax dark brown with a broad white stripe on each side. Abdomen dark brown with a bluish lustre, and broad white band at the hind edge of the second segment. Anal tuft dark brown with two white streaks. Underside of body wholly brown, also the legs. Anal tuft with two short white streaks, not reaching the end. Fore wings opaque, brown with a purplish lustre, and a small white spot composed of scales at the end of the cell. Hind wings opaque, wholly brown. Underside of wings same as above, but the fore wings are marked with whitish along the costal region. Expanse, 28–30 mm.



Habitat.—Phoenix, Arizona.

Described from two males. Types, Collection Am. Mus. Nat. Hist., and Charles Palm. The figure is enlarged one fourth.